

RATIONAL AND TRIGONOMETRIC DEGENERATION OF THE DOUBLE AFFINE HECKE ALGEBRA OF TYPE A

TAKESHI SUZUKI

ABSTRACT. We study a connection between the representation theory of the rational Cherednik algebra of type GL_n and the representation theory of the degenerate double affine Hecke algebra (the degenerate DAHA). We focus on an algebra embedding from the rational Cherednik algebra to the degenerate DAHA and investigate the induction functor through this embedding. We prove that this functor embeds the category $\mathcal{O}$ for the rational Cherednik algebra fully faithfully into the category $\mathcal{O}$ for the degenerate DAHA. We also study the full subcategory $\mathcal{O}^{ss}$ of $\mathcal{O}$ consisting of those modules which are semisimple with respect to the commutative subalgebra generated by Cherednik-Dunkl operators. A classification of all irreducible modules in $\mathcal{O}^{ss}$ for the rational Cherednik algebra is obtained from the corresponding result for the degenerate DAHA.

Introduction

We study a connection between the representation theories of two algebras of type GL_n ; the rational Cherednik algebra $\mathbf{H}_\kappa$ and the degenerate double affine Hecke algebra (or the trigonometric Cherednik algebra) $\tilde{\mathbf{H}}_\kappa$. These two algebras can be thought (see [EG]) as a rational and trigonometric degeneration of the double affine Hecke algebra introduced by Cherednik [Ch1, Ch2] respectively.

The categories of finite dimensional representations for the rational and trigonometric Cherednik algebras (of type SL_n) have been related through deformation argument ([BEG, Proposition 7.1][Ch3]).

In this paper, we relate what is called the categories $\mathcal{O}$ ([DO, GGOR, VV]) for $\mathbf{H}_\kappa$ and $\tilde{\mathbf{H}}_\kappa$ by a different approach. We give an algebra embedding from $\mathbf{H}_\kappa$ to $\tilde{\mathbf{H}}_\kappa$, which extends to an algebra isomorphism from a localization of $\mathbf{H}_\kappa$ to $\tilde{\mathbf{H}}_\kappa$ (Proposition 3.1), and investigate the induction functor $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (-) : \mathbf{H}_\kappa\text{-Mod} \rightarrow \tilde{\mathbf{H}}_\kappa\text{-Mod}$ through this embedding.

By restricting this functor, we obtain the functor from $\mathcal{O}(\mathbf{H}_\kappa)$ into $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$, where $\mathcal{O}(\mathbf{H}_\kappa)$ and $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ denote the category $\mathcal{O}$ for $\mathbf{H}_\kappa$ and

$\tilde{\mathbf{H}}_\kappa$ respectively. The main result of this paper (Theorem 5.4) asserts that this functor embeds $\mathcal{O}(\mathbf{H}_\kappa)$ into $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ fully faithfully. Moreover it follows that our functor sends a standard module of $\mathbf{H}_\kappa$ to a standard module of $\tilde{\mathbf{H}}_\kappa$, and an irreducible module to an irreducible module.

Through this functor, one can transform results on the one side to the other. For example, the multiplicity of an irreducible module in the composition series of a standard module for $\mathbf{H}_\kappa$ is equal to the corresponding multiplicity for $\tilde{\mathbf{H}}_\kappa$. Note that $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ has been also related to the category $\mathcal{O}$ for the (original) double affine Hecke algebra ([VV]), and hence this multiplicity is also equal to the corresponding multiplicity for the double affine Hecke algebra, which can be expressed in terms of Kazhdan-Lusztig polynomials as shown in [Va] by a geometric method.

We also focus on the full subcategory $\mathcal{O}^{ss}$ of $\mathcal{O}$ consisting of those modules which are semisimple with respect to the commutative subalgebra which is realized as the algebra generated by Cherednik-Dunkl operators on the polynomial representation. This class of representations for $\tilde{\mathbf{H}}_\kappa$ have been studied in [Ch3, SV] (see also [Ch4]), and in particular the classification of all irreducible modules in $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$ has been obtained. It turns out that the induction functor transforms $\mathcal{O}^{ss}(\mathbf{H}_\kappa)$ into $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$, and hence the classification (Theorem 7.2) of the irreducible modules in $\mathcal{O}^{ss}(\mathbf{H}_\kappa)$ follows from the classification result for $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$.

Acknowledgments. The author would like to thank Prof. Opdam for discussion and his lecture on Cherednik algebras while his stay in RIMS. The author is also very grateful to Prof. Kashiwara for important suggestion and comments.

1. THE AFFINE WEYL GROUP

Let $\mathbb{F}$ be an algebraically closed field of characteristic 0. Let $\mathbb{Q}$ and $\mathbb{Z}$ denote the set of rational numbers and the set of integers respectively, which are subrings of $\mathbb{F}$.

Throughout this paper, we use the notation

$$[a, b] = \{a, a + 1, \dots, b\} \subset \mathbb{F}$$

for $a, b \in \mathbb{F}$ such that $b - a \in \mathbb{Z}$.

Let $n \in \mathbb{Z}_{\geq 2}$. Let $\mathfrak{S}_n$ denote the symmetric group of degree n . For $i, j \in [1, n]$ with $i \neq j$, let s_{ij} denote the transposition of i and j . In particular, we write s_i ($i \in [1, n - 1]$) for the simple reflection which transposes i and $i + 1$.

Let $P = \bigoplus_{i \in [1, n]} \mathbb{Z} \epsilon_i$ denote the free $\mathbb{Z}$ -module generated by $\epsilon_1, \epsilon_2, \dots, \epsilon_n$, on which $\mathfrak{S}_n$ acts naturally. Define the extended affine Weyl group $\widehat{\mathfrak{S}}_n$ as the semidirect product $P \rtimes \mathfrak{S}_n$.

Put $\pi = t_{\epsilon_1} s_1 s_2 \cdots s_{n-1}$ and $s_0 = t_{\epsilon_1 - \epsilon_n} s_{1n}$, where t_η denotes the element of $\widehat{\mathfrak{S}}_n$ corresponding to $\eta \in P$. Then $\{\pi, s_0, s_1, \dots, s_{n-1}\}$ is a set of generators of the group $\widehat{\mathfrak{S}}_n$. We put $P_\circ = \bigoplus_{i \in [1, n]} \mathbb{Z}_{\geq 0} \epsilon_i \subset P$ and $\widehat{\mathfrak{S}}_n^\circ = P_\circ \rtimes \mathfrak{S}_n \subset \widehat{\mathfrak{S}}_n$. Observe that $\widehat{\mathfrak{S}}_n^\circ$ is the semigroup (with unit) generated by $\{\pi, s_1, s_2, \dots, s_{n-1}\}$.

For a semigroup G , let $\mathbb{F}G$ denote the semigroup algebra of G .

Lemma 1.1. *The linear map $\mathbb{F}[\pi, \pi^{-1}] \otimes \mathbb{F}\widehat{\mathfrak{S}}_n^\circ \rightarrow \mathbb{F}\widehat{\mathfrak{S}}_n$ given by $\pi^k \otimes w \mapsto \pi^k w$ induces a linear isomorphism $\mathbb{F}[\pi, \pi^{-1}] \otimes_{\mathbb{F}[\pi]} \mathbb{F}\widehat{\mathfrak{S}}_n^\circ \xrightarrow{\sim} \mathbb{F}\widehat{\mathfrak{S}}_n$.*

Proof. Let $w \in \widehat{\mathfrak{S}}_n$. Note that $\pi^n = t_{\epsilon_1 + \epsilon_2 + \dots + \epsilon_n}$. Hence there exist some $k \in \mathbb{Z}$ and $\bar{w} \in \widehat{\mathfrak{S}}_n^\circ$ such that $w = \pi^k \bar{w}$. It follows that the correspondence $w \mapsto \pi^k \otimes \bar{w}$ gives rise to a well-defined linear map $\mathbb{F}\widehat{\mathfrak{S}}_n \rightarrow \mathbb{F}[\pi, \pi^{-1}] \otimes_{\mathbb{F}[\pi]} \mathbb{F}\widehat{\mathfrak{S}}_n^\circ$, and it gives the inverse of the induced map $\mathbb{F}[\pi, \pi^{-1}] \otimes_{\mathbb{F}[\pi]} \mathbb{F}\widehat{\mathfrak{S}}_n^\circ \rightarrow \mathbb{F}\widehat{\mathfrak{S}}_n$. $\square$

We denote the polynomial ring and the Laurent polynomial ring in n -variables $x_1, x_2, \dots, x_n$ by $\mathbb{F}[\underline{x}]$ and $\mathbb{F}[\underline{x}^{\pm 1}]$ respectively:

$$\mathbb{F}[\underline{x}] = \mathbb{F}[x_1, x_2, \dots, x_n], \quad \mathbb{F}[\underline{x}^{\pm 1}] = \mathbb{F}[x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_n^{\pm 1}].$$

In the sequel, we identify $\mathbb{F}P$ with $\mathbb{F}[\underline{x}^{\pm 1}]$, and identify $\mathbb{F}\widehat{\mathfrak{S}}_n$ with $\mathbb{F}[\underline{x}^{\pm 1}] \rtimes \mathfrak{S}_n$ as algebras through the correspondence $t_{\epsilon_i} \mapsto x_i$ ($i \in [1, n]$). Observe that $\mathbb{F}\widehat{\mathfrak{S}}_n^\circ$ is identified with $\mathbb{F}[\underline{x}] \rtimes \mathfrak{S}_n$.

2. CHEREDNIK ALGEBRAS

Fix $\kappa \in \mathbb{F} \setminus \{0\}$.

Let $\mathcal{F}$ denote the associative algebra generated freely by the algebras $\mathbb{F}\widehat{\mathfrak{S}}_n^\circ = \mathbb{F}[\underline{x}] \rtimes \mathfrak{S}_n$ and $\mathbb{F}[\underline{y}] = \mathbb{F}[y_1, y_2, \dots, y_n]$ such that the natural inclusions $\mathbb{F}\widehat{\mathfrak{S}}_n^\circ \rightarrow \mathcal{F}$ and $\mathbb{F}[\underline{y}] \rightarrow \mathcal{F}$ are algebra homomorphisms. The rational Cherednik algebra $\mathbf{H}_\kappa$ of type GL_n is the unital associative $\mathbb{F}$ -algebra defined as the quotient algebra of $\mathcal{F}$ by the following relations:

$$\begin{aligned} s_i y_j &= y_{s_i(j)} s_i \quad (i \in [1, n-1], j \in [1, n]), \\ [y_i, x_j] &= \begin{cases} \kappa + \sum_{i \neq k} s_{ik} & (i = j), \\ -s_{ij} & (i \neq j). \end{cases} \end{aligned}$$

Let $\widetilde{\mathcal{F}}$ denote the algebra generated freely by the algebras $\mathbb{F}\widehat{\mathfrak{S}}_n = \mathbb{F}[\underline{x}^{\pm 1}] \rtimes \mathfrak{S}_n$ and $\mathbb{F}[\underline{u}] = \mathbb{F}[u_1, u_2, \dots, u_n]$ such that the natural inclusions $\mathbb{F}\widehat{\mathfrak{S}}_n \rightarrow \widetilde{\mathcal{F}}$ and $\mathbb{F}[\underline{u}] \rightarrow \widetilde{\mathcal{F}}$ are algebra homomorphisms. The

degenerate double affine Hecke algebra (or the trigonometric Cherednik algebra) $\tilde{\mathbf{H}}_\kappa$ of type GL_n is the unital associative $\mathbb{F}$ -algebra defined as the quotient algebra of $\tilde{\mathcal{F}}$ by the following relations:

$$\begin{aligned} s_i u_i &= u_{i+1} s_i - 1, \quad s_i u_{i+1} = u_i s_i + 1 \quad (i \in [1, n-1]), \\ s_i u_j &= u_j s_i \quad (i \in [1, n-1], j \in [1, n], j \neq i, i+1), \\ [u_i, x_j] &= \begin{cases} \kappa x_i + \sum_{1 \leq k < i} x_k s_{ki} + \sum_{i < k \leq n} x_i s_{ik} & (i = j), \\ -x_j s_{ji} & (i > j), \\ -x_i s_{ij} & (i < j). \end{cases} \end{aligned}$$

Remark 2.1. The degenerate double affine Hecke algebra can be defined in an alternative way in terms of the generators

$$\{\pi, \pi^{-1}, s_0, \dots, s_{n-1}, u_1, \dots, u_n\}.$$

It is well-known that the two definitions coincide (see e.g. [AST, Proposition 1.3.7]). Note in particular that the following relations hold:

$$(2.1) \quad \pi u_i = u_{i+1} \pi \quad (i \in [1, n-1]), \quad \pi u_n = (u_1 - \kappa) \pi.$$

Proposition 2.2. (Cherednik, [EG, Theorem 1.3])

(i) *Multiplication in $\mathbf{H}_\kappa$ induces a linear isomorphism*

$$\mathbb{F}[\underline{x}] \otimes \mathbb{F}\mathfrak{S}_n \otimes \mathbb{F}[\underline{y}] \xrightarrow{\sim} \mathbf{H}_\kappa.$$

(ii) *Multiplication in $\tilde{\mathbf{H}}_\kappa$ induces a linear isomorphism*

$$\mathbb{F}[\underline{x}^{\pm 1}] \otimes \mathbb{F}\mathfrak{S}_n \otimes \mathbb{F}[\underline{u}] \xrightarrow{\sim} \tilde{\mathbf{H}}_\kappa.$$

The subalgebra $\mathbf{H}^{\text{aff}} = \mathbb{F}\mathfrak{S}_n \cdot \mathbb{F}[\underline{u}]$ of $\tilde{\mathbf{H}}_\kappa$ is called *the degenerate affine Hecke algebra* of type GL_n .

3. THE ALGEBRA EMBEDDING AND THE INDUCTION FUNCTOR

The algebra structure of $\mathbf{H}_\kappa$ is extended to the algebra structure of $\mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa = \mathbb{F}[\underline{x}^{\pm 1}] \otimes \mathbb{F}\mathfrak{S}_n \otimes \mathbb{F}[\underline{y}]$.

Proposition 3.1. (i) *There exists a unique algebra homomorphism $\iota : \mathbf{H}_\kappa \rightarrow \tilde{\mathbf{H}}_\kappa$ such that*

$$\begin{aligned} \iota(w) &= w \quad (w \in \mathfrak{S}_n), \\ \iota(x_i) &= x_i \quad (i \in [1, n]), \\ \iota(y_i) &= x_i^{-1} \left(u_i - \sum_{1 \leq j < i} s_{ji} \right) \quad (i \in [1, n]), \end{aligned}$$

and moreover it is injective.

(ii) The algebra embedding $\iota : \mathbf{H}_\kappa \rightarrow \tilde{\mathbf{H}}_\kappa$ is extended to the algebra isomorphism $\mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa \xrightarrow{\sim} \tilde{\mathbf{H}}_\kappa$, whose inverse map $j : \tilde{\mathbf{H}}_\kappa \rightarrow \mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa$ is defined by

$$\begin{aligned} j(w) &= w \quad (w \in \mathfrak{S}_n), \\ j(x_i) &= x_i \quad (i \in [1, n]), \\ j(u_i) &= x_i y_i + \sum_{1 \leq j < i} s_{ji} \quad (i \in [1, n]). \end{aligned}$$

Proof. By checking the defining relations of $\mathbf{H}_\kappa$ (resp. $\tilde{\mathbf{H}}_\kappa$) by direct calculations, one can show that the assignment ι (resp. j) are uniquely extended to the algebra homomorphism $\mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa \rightarrow \tilde{\mathbf{H}}_\kappa$ (resp. $\tilde{\mathbf{H}}_\kappa \rightarrow \mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa$). It is easy to see that $j \circ \iota = \text{id}_{\tilde{\mathbf{H}}_\kappa}$ and $\iota \circ j = \text{id}_{\mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa}$. This proves (ii). Now (i) follows immediately because the natural map $\mathbf{H}_\kappa \rightarrow \mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa$ is injective by Proposition 2.2-(i). $\square$

Remark 3.2. Recall the polynomial representation $\mathbf{H}_\kappa \rightarrow \text{End}_{\mathbb{F}}(\mathbb{F}[\underline{x}])$, where the Dunkl operators

$$Y_i = \kappa \frac{\partial}{\partial x_i} + \sum_{j \neq i} \frac{1}{x_i - x_j} (1 - s_{ij}) \quad (i \in [1, n])$$

appear as the images of y_i ($i \in [1, n]$). The operators $U_i := x_i Y_i + \sum_{1 \leq j < i} s_{ji}$ ($i \in [1, n]$), which are also pairwise commutative, are introduced by Cherednik ([Ch1]) as a trigonometric analogue of the Dunkl operators. The original and trigonometric Dunkl operators play important role in the theory of orthogonal polynomials (see e.g. [DX]). Note in particular that the simultaneous eigenvectors of $\mathbb{F}[\underline{x}]$ with respect to the trigonometric Dunkl operators $U_1, \dots, U_n$ are the nonsymmetric Jack polynomials.

Note also that generalization of trigonometric Dunkl operators to other types than A has been studied ([Ka, DO]).

Remark 3.3. The restriction of the isomorphism j to the subalgebra $\mathbf{H}^{\text{aff}}$ gives an algebra embedding $\mathbf{H}^{\text{aff}} \rightarrow \mathbf{H}_\kappa$. More generally, it was shown in [EG, Proposition 4.3] that the assignment

$$u_i \mapsto ax_i + by_i + x_i y_i + \sum_{1 \leq j < i} s_{ji} \quad (i \in [1, n]), \quad w \mapsto w \quad (w \in \mathfrak{S}_n)$$

extends to an algebra embedding $\mathbf{H}^{\text{aff}} \rightarrow \mathbf{H}_\kappa$ for any $a, b \in \mathbb{F}$. When $b = 0$, this embedding extends to the algebra isomorphism $j_a : \tilde{\mathbf{H}}_\kappa \rightarrow \mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa$ such that $j_a(x_i) = x_i$ ($i \in [1, n]$).

In the rest of the paper, we often identify $\tilde{\mathbf{H}}_\kappa$ with $\mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa$, and regard $\mathbf{H}_\kappa$ as a subalgebra of $\tilde{\mathbf{H}}_\kappa$.

Let us consider the induction functor $M \mapsto \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ from the category $\mathbf{H}_\kappa\text{-Mod}$ of $\mathbf{H}_\kappa$ -modules to the category $\tilde{\mathbf{H}}_\kappa\text{-Mod}$ of $\tilde{\mathbf{H}}_\kappa$ -modules. Since $\tilde{\mathbf{H}}_\kappa$ can be seen as a localization of $\mathbf{H}_\kappa$ by Proposition 3.1, we have

Corollary 3.4. *The $\tilde{\mathbf{H}}_\kappa$ is flat as an $\mathbf{H}_\kappa$ -module, or equivalently, the induction functor $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (-) : \mathbf{H}_\kappa\text{-Mod} \rightarrow \tilde{\mathbf{H}}_\kappa\text{-Mod}$ is exact.*

Corollary 3.5. *Let M be an irreducible $\mathbf{H}_\kappa$ -module. Then $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is an irreducible $\tilde{\mathbf{H}}_\kappa$ -module or zero.*

Proof. Follows easily from the isomorphism $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M \cong \mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} M$. $\square$

For later use, we give another expression of $\tilde{\mathbf{H}}_\kappa$, which follows from Lemma 1.1.

Corollary 3.6. *The linear map $\mathbb{F}[\pi, \pi^{-1}] \otimes \mathbf{H}_\kappa \rightarrow \tilde{\mathbf{H}}_\kappa$ given by $\pi^k \otimes v \mapsto \pi^k v$, induces a linear isomorphism $\mathbb{F}[\pi, \pi^{-1}] \otimes_{\mathbb{F}[\pi]} \mathbf{H}_\kappa \xrightarrow{\sim} \tilde{\mathbf{H}}_\kappa$.*

We will use the following lemma later.

Lemma 3.7. *There exists an algebra isomorphism $\sigma : \tilde{\mathbf{H}}_\kappa \rightarrow \tilde{\mathbf{H}}_{-\kappa}$ such that*

$$\sigma(s_i) = -s_i \ (i \in [0, n-1]), \ \sigma(x_i) = (-1)^{n-1} x_i, \ \sigma(u_i) = -u_i \ (i \in [1, n]),$$

and its restriction to $\mathbf{H}_\kappa$ gives an isomorphism $\sigma : \mathbf{H}_\kappa \xrightarrow{\sim} \mathbf{H}_{-\kappa}$.

4. THE CATEGORY $\mathcal{O}$

Put $\mathbb{F}[\underline{y}]_+ = \sum_{i \in [1, n]} y_i \mathbb{F}[\underline{y}] \subset \mathbf{H}_\kappa$. An element v of an $\mathbf{H}_\kappa$ -module M is said to be $\mathbb{F}[\underline{y}]_+$ -nilpotent if there exists $k > 0$ such that $y_i^k v = 0$ for all $i \in [1, n]$. An $\mathbf{H}_\kappa$ -module M is said to be locally nilpotent for $\mathbb{F}[\underline{y}]_+$, if any element of M is $\mathbb{F}[\underline{y}]_+$ -nilpotent. Let $\mathcal{O}(\mathbf{H}_\kappa)$ denote the full subcategory of $\mathbf{H}_\kappa\text{-Mod}$ consisting of those modules which are finitely generated over $\mathbf{H}_\kappa$ and locally nilpotent for $\mathbb{F}[\underline{y}]_+$.

Let $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ denote the full subcategory of $\tilde{\mathbf{H}}_\kappa\text{-Mod}$ consisting of those modules which are finitely generated over $\tilde{\mathbf{H}}_\kappa$ and locally finite for $\mathbb{F}[\underline{u}]$, i.e., a finitely generated $\tilde{\mathbf{H}}_\kappa$ -module M is in $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ if $\mathbb{F}[\underline{u}]v < \infty$ for any $v \in M$.

Proposition 4.1. (i) ([GGOR, Corollary 2.16]) *The category $\mathcal{O}(\mathbf{H}_\kappa)$ is a Serre subcategory of $\mathbf{H}_\kappa\text{-Mod}$, and any object in $\mathcal{O}(\mathbf{H}_\kappa)$ has a finite composition series.*

(ii) ([VV, Proposition 2.1, Proposition 2.2]) *The category $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ is a Serre subcategory of $\tilde{\mathbf{H}}_\kappa\text{-Mod}$, and any object in $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ has a finite composition series.*

For $d \in \mathbb{Z}_{\geq 0}$, define $\mathbb{F}[\underline{x}]_{(d)}$ as the subspace of $\mathbb{F}[\underline{x}]$ spanned by $\{x_1^{d_1}x_2^{d_2}\cdots x_n^{d_n} \mid \sum_{i \in [1,n]} d_i = d\}$. We have $\bigoplus_{d=0}^\infty \mathbb{F}[\underline{x}]_{(d)} = \mathbb{F}[\underline{x}]$. Define $\mathbb{F}[\underline{y}]_{(d)}$ similarly.

Proposition 4.2. *Let M be an object of $\mathcal{O}(\mathbf{H}_\kappa)$. Then $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is an object of $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$.*

Proof. Recall that, under the identification $\tilde{\mathbf{H}}_\kappa \cong \mathbb{F}[\underline{x}^{\pm 1}] \otimes_{\mathbb{F}[\underline{x}]} \mathbf{H}_\kappa$, the element $u_i \in \tilde{\mathbf{H}}_\kappa$ is expressed as $x_i y_i + \sum_{j < i} s_{ji}$, and it is contained in $\mathbf{H}_\kappa \subset \tilde{\mathbf{H}}_\kappa$.

Let M be an object of $\mathcal{O}(\mathbf{H}_\kappa)$ and take $v \in M$. As a step, we will prove that $\mathbb{F}[\underline{u}]v \subseteq M$ is finite-dimensional. It is easy to see that

$$u_i^d = \left(x_i y_i + \sum_{j < i} s_{ji} \right)^d \in \bigoplus_{k \leq d} \mathbb{F}[\underline{x}]_{(k)} \cdot \mathbb{F}\mathfrak{S}_n \cdot \mathbb{F}[\underline{y}]_{(k)}.$$

Since M is locally nilpotent for $\mathbb{F}[\underline{y}]_+$, there exists $K \in \mathbb{Z}_{\geq 0}$ such that $\mathbb{F}[\underline{y}]_{(k)}v = 0$ if $k > K$. Hence for any $i \in [1, n]$ and $d \in \mathbb{Z}_{\geq 0}$, it holds that $u_i^d v$ is contained in $\bigoplus_{k \leq K} \mathbb{F}[\underline{x}]_{(k)} \cdot \mathbb{F}\mathfrak{S}_n \cdot \mathbb{F}[\underline{y}]_{(k)}v$, which is finite-dimensional. This implies $\dim(\mathbb{F}[\underline{u}]v) < \infty$.

By Corollary 3.6, any element in $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is of the form $\pi^k \bar{v}$ for some $k \in \mathbb{Z}_{\leq 0}$ and $v \in M$, where $\bar{v}$ denotes the image of $1 \otimes v$ in $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$. Since $\pi \mathbb{F}[\underline{u}] \pi^{-1} = \mathbb{F}[\underline{u}]$, we have $\dim(\mathbb{F}[\underline{u}] \pi^k \bar{v}) = \dim(\mathbb{F}[\underline{u}]v) < \infty$. $\square$

Let $\text{Res}_{\mathbf{H}_\kappa}(-)$ denote the restriction functor $\tilde{\mathbf{H}}_\kappa\text{-Mod} \rightarrow \mathbf{H}_\kappa\text{-Mod}$.

Lemma 4.3. (i) *Let M be an object of $\mathcal{O}(\mathbf{H}_\kappa)$. Then the natural embedding $M \hookrightarrow \tilde{\mathbf{H}}_\kappa \otimes M$ induces an injective $\mathbf{H}_\kappa$ -homomorphism*

$$M \hookrightarrow \text{Res}_{\mathbf{H}_\kappa} \left(\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M \right).$$

(ii) *Let N be an object of $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$. Then the product map $\tilde{\mathbf{H}}_\kappa \otimes N \rightarrow N$ induces an $\tilde{\mathbf{H}}_\kappa$ -isomorphism*

$$\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (\text{Res}_{\mathbf{H}_\kappa} N) \xrightarrow{\sim} N.$$

Proof. (i) It is obvious that the map is an $\mathbf{H}_\kappa$ -homomorphism. Let us prove that it is injective. Let $v \in M \setminus \{0\}$. Put $\omega = \sum_{i \in [1,n]} y_i \in \mathbb{F}[\underline{y}]_{(1)} \subset \mathbf{H}_\kappa$. Then we have $[\omega^p, x_i] = p\kappa\omega^{p-1}$ for any $p \in \mathbb{Z}_{\geq 1}$. Since M is locally nilpotent for $\mathbb{F}[\underline{y}]_+$, there exists $p_0 \in \mathbb{Z}_{\geq 1}$ such that $\omega^{p_0-1}v \neq 0$

and $\omega^{p_0}v = 0$. For any $i \in [1, n]$, we have $\omega^{p_0}x_i v = p_0 \kappa \omega^{p_0-1}v \neq 0$, and hence $x_i v \neq 0$. This implies $fv \neq 0$ for any monomial f in $x_1, \dots, x_n$. This proves the statement.

(ii) Follows easily from the isomorphism $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} N \cong \mathbb{F}[\pi^{\pm 1}] \otimes_{\mathbb{F}[\pi]} N$. $\square$

5. THE ADJOINT FUNCTOR $\mathfrak{N}$

Now we introduce a functor $\mathcal{O}(\tilde{\mathbf{H}}_\kappa) \rightarrow \mathcal{O}(\mathbf{H}_\kappa)$, which turns out to be the right adjoint functor of $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (-)$.

For an $\tilde{\mathbf{H}}_\kappa$ -module M , denote by $\mathfrak{N}(M)$ the subspace of M consisting of the $\mathbb{F}[y]_+$ -nilpotent elements.

Lemma 5.1. *If M is in $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$, then $\mathfrak{N}(M)$ is an $\mathbf{H}_\kappa$ -submodule of $\text{Res}_{\mathbf{H}_\kappa} M$ and it is finitely generated over $\mathbf{H}_\kappa$.*

Proof. Let $v \in \mathfrak{N}(M)$. Then it is obvious that $s_i v$ and $y_i v$ are in $\mathfrak{N}(M)$. It is easily shown by induction that $\mathbb{F}[y]_{(p)} x_i \subseteq \mathbb{F}[\underline{x}]_{(1)} \mathbb{F}[y]_{(p)} + \mathbb{F}\mathfrak{S}_n \mathbb{F}[y]_{(p-1)}$ for any $p \in \mathbb{Z}_{\geq 1}$. This implies $x_i v \in \mathfrak{N}(M)$ and hence $\mathfrak{N}(M)$ is an $\mathbf{H}_\kappa$ -submodule.

Let us prove that $\mathfrak{N}(M)$ is finitely generated. Take a sequence

$$(5.1) \quad 0 = N^{(0)} \subsetneq N^{(1)} \subsetneq \dots \subsetneq N^{(k)} \subsetneq \dots$$

of finitely generated $\mathbf{H}_\kappa$ -submodules of $\mathfrak{N}(M)$. Since each $N^{(k)}$ is in $\mathcal{O}(\mathbf{H}_\kappa)$, it follows from Lemma 4.3-(i) that $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} N^{(k)} \subsetneq \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} N^{(k+1)}$ for all k . By the exactness of the induction functor on $\mathbf{H}_\kappa\text{-Mod}$ and by Lemma 4.3-(ii), we have $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} \mathfrak{N}(M) \subseteq \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} \text{Res}_{\mathbf{H}_\kappa}(M) \cong M$. Hence $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} \mathfrak{N}(M)$ is in $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ and its composition series is finite. Therefore the sequence (5.1) must end in finite steps, and we have $N^{(K)} = \mathfrak{N}(M)$ for some $K \in \mathbb{Z}_{\geq 0}$. $\square$

Therefore we obtain the functor $\mathfrak{N} : \mathcal{O}(\tilde{\mathbf{H}}_\kappa) \rightarrow \mathcal{O}(\mathbf{H}_\kappa)$.

Corollary 5.2. *The functor $\mathfrak{N} : \mathcal{O}(\tilde{\mathbf{H}}_\kappa) \rightarrow \mathcal{O}(\mathbf{H}_\kappa)$ is the right adjoint functor of the functor $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (-) : \mathcal{O}(\mathbf{H}_\kappa) \rightarrow \mathcal{O}(\tilde{\mathbf{H}}_\kappa)$; namely, we have a natural isomorphism*

$$(5.2) \quad \text{Hom}_{\tilde{\mathbf{H}}_\kappa}(\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M, N) \cong \text{Hom}_{\mathbf{H}_\kappa}(M, \mathfrak{N}(N))$$

for any M in $\mathcal{O}(\mathbf{H}_\kappa)$ and any N in $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$.

Proposition 5.3. *Let M be in $\mathcal{O}(\mathbf{H}_\kappa)$. Then there exists a natural $\tilde{\mathbf{H}}_\kappa$ -isomorphism $\mathfrak{N}(\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M) \cong M$.*

Proof. Using Lemma 4.3-(i), we have the following sequence of embeddings of $\mathbf{H}_\kappa$ -modules:

$$(5.3) \quad M \xhookrightarrow{i} \mathfrak{N}(\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M) \hookrightarrow \text{Res}_{\mathbf{H}_\kappa}(\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M).$$

By applying $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (-)$, we have embeddings of $\tilde{\mathbf{H}}_\kappa$ -modules

$$(5.4) \quad \tilde{M} \xhookrightarrow{\tilde{i}} \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (\mathfrak{N}(\tilde{M})) \hookrightarrow \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (\text{Res}_{\mathbf{H}_\kappa} \tilde{M}),$$

where we put $\tilde{M} = \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$. By combining it with the isomorphism $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (\text{Res}_{\mathbf{H}_\kappa} \tilde{M}) \xrightarrow{\sim} \tilde{M}$ given in Lemma 4.3-(ii), we have an $\tilde{\mathbf{H}}_\kappa$ -homomorphism $\tilde{M} \rightarrow \tilde{M}$, and it easily follows that this map equals the identity map. Therefore the embedding $\tilde{i}$ in (5.4) is an isomorphism, and hence we have $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (\text{Coker } i) = \text{Coker } \tilde{i} = 0$. This implies $\text{Coker } i = 0$ by Lemma 4.3-(i). Namely, the embedding $i : M \rightarrow \mathfrak{N}(\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M)$ in (5.3) is an isomorphism. $\square$

Theorem 5.4. *The functor $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (-) : \mathcal{O}(\mathbf{H}_\kappa) \rightarrow \mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ is exact and fully faithful.*

Proof. The exactness has been shown in Corollary 3.4. The claim that $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} (-)$ is fully faithful follows from Corollary 5.2 and Proposition 5.3 $\square$

Remark 5.5. It is conjectured in [GGOR, Remark 5.17] that the category $\mathcal{O}(\mathbf{H}_\kappa)$ is equivalent to the category of finite dimensional modules over the q -Schur algebra with $q = \exp(\frac{2\pi i}{\kappa})$. The analogous statement for $\tilde{\mathbf{H}}_\kappa$ has been proved in [VV, Theorem 5.2], where they gave a categorical equivalence between $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ and the category of finite dimensional modules over the affine q -Schur algebra.

6. STANDARD AND IRREDUCIBLE MODULES

Introduce the subalgebra $\mathbf{H}_{\geq 0} = \mathbb{F}\mathfrak{S}_n \cdot \mathbb{F}[\underline{y}]$ of $\mathbf{H}_\kappa$. Let $\phi : \mathbf{H}_{\geq 0} \rightarrow \mathbb{F}\mathfrak{S}_n$ be the algebra homomorphism given by

$$\phi(s_i) = s_i \ (i \in [1, n-1]), \quad \phi(x_i) = 0 \ (i \in [1, n]).$$

For a finite-dimensional $\mathfrak{S}_n$ -module E , let ${}^\phi E$ denote the $\mathbf{H}_{\geq 0}$ -modules given by the composition $\mathbf{H}_{\geq 0} \xrightarrow{\phi} \mathbb{F}\mathfrak{S}_n \rightarrow \text{End}(E)$.

We define the standard module of $\mathbf{H}_\kappa$ associated with E by

$$\Delta_\kappa(E) = \mathbf{H}_\kappa \otimes_{\mathbf{H}_{\geq 0}} {}^\phi E.$$

There exists an algebra homomorphism $\psi : \mathbf{H}^{\text{aff}} \rightarrow \mathbb{F}\mathfrak{S}_n$ given by

$$\psi(s_i) = s_i \ (i \in [1, n-1]), \quad \psi(u_i) = \sum_{j < i} s_{ji} \ (i \in [1, n]),$$

and define the $\mathbf{H}^{\text{aff}}$ -module ${}^\psi E$ similarly as ${}^\phi E$. We define the standard module of $\tilde{\mathbf{H}}_\kappa$ associated with E by

$$\tilde{\Delta}_\kappa(E) = \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}^{\text{aff}}} {}^\psi E.$$

Observe that the standard modules $\Delta_\kappa(E)$ (resp. $\tilde{\Delta}_\kappa(E)$) are objects of $\mathcal{O}(\mathbf{H}_\kappa)$ (resp. $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$).

Theorem 6.1. (i) *Let E be a finite-dimensional $\mathfrak{S}_n$ -module. Then*

$$\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} \Delta_\kappa(E) \cong \tilde{\Delta}_\kappa(E).$$

(ii) *Let M be in $\mathcal{O}(\mathbf{H}_\kappa)$. Then $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is irreducible if and only if M is irreducible.*

Proof. (i) Observe that y_i acts trivially on the image of ${}^\psi E$ in $\tilde{\Delta}_\kappa(E)$. Thus we have an $\mathbf{H}_\kappa$ -homomorphism $\Delta_\kappa(E) \rightarrow \tilde{\Delta}_\kappa(E)$, and hence a surjective $\tilde{\mathbf{H}}_\kappa$ -homomorphism $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} \Delta_\kappa(E) \rightarrow \tilde{\Delta}_\kappa(E)$. This is isomorphism since both sides are isomorphic to $\mathbb{F}P \otimes E$ as $\mathbb{F}P$ -modules. (ii) The statement follows immediately from Theorem 5.4. $\square$

Let E be a finite-dimensional irreducible $\mathfrak{S}_n$ -module. A similar argument as in the theory of highest weight modules for Lie algebras implies that the standard module $\Delta_\kappa(E)$ has a unique simple quotient, and any simple object in $\mathcal{O}(\mathbf{H}_\kappa)$ is given as such a quotient [GGOR, Proposition 2.11].

Corollary 6.2. (*cf.* [AST, Proposition 2.5.3] [Su, Theorem 8.2]) *Let E be a finite-dimensional irreducible $\mathfrak{S}_n$ -module. Then $\tilde{\Delta}_\kappa(E)$ has a unique simple quotient module.*

We denote by $\mathcal{L}_\kappa(E)$ and $\tilde{\mathcal{L}}_\kappa(E)$ the unique simple quotient of $\Delta_\kappa(E)$ and $\tilde{\Delta}_\kappa(E)$ respectively. As a direct consequence of Theorem 6.1, we have $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} \mathcal{L}_\kappa(E) \cong \tilde{\mathcal{L}}_\kappa(E)$,

Corollary 6.3. *Let E and F be finite-dimensional irreducible $\mathfrak{S}_n$ -modules. Then*

$$[\Delta_\kappa(E) : \mathcal{L}_\kappa(F)] = [\tilde{\Delta}_\kappa(E) : \tilde{\mathcal{L}}_\kappa(F)],$$

where $[M : N]$ denotes the multiplicity of the irreducible module N in the composition series of M .

Remark 6.4. The category $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ has been also related to the category $\mathcal{O}$ for the (original) double affine Hecke algebra ([VV, Proposition 2.5]), and, as a consequence, the multiplicity for $\tilde{\mathbf{H}}_\kappa$ in Corollary 6.3 equals the corresponding multiplicity for the double affine Hecke algebra. The

latter can be expressed in terms of the Kazhdan-Lusztig polynomials ([Va, Theorem 8.5]).

7. $\mathbb{F}[\underline{u}]$ -SEMISIMPLE MODULES

Recall that $\mathbf{H}_\kappa$ includes the degenerate affine Hecke algebra $\mathbf{H}^{\text{aff}}$ and in particular the commutative subalgebra $\mathbb{F}[\underline{u}]$.

For an $\mathbf{H}^{\text{aff}}$ -module M and $\zeta = (\zeta_1, \dots, \zeta_n) \in \mathbb{F}^n$, set

$$M_\zeta = \{v \in M \mid u_i v = \zeta_i v \text{ for all } i \in [1, n]\}.$$

We call an element of M_ζ a weight vector of weight ζ .

Define the category $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$ as the full subcategory of $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ consisting of $\mathbb{F}[\underline{u}]$ -semisimple modules, i.e., an object M in $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$ is in $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$ if $M = \bigoplus_{\zeta \in \mathbb{F}^n} M_\zeta$. Similarly, define the category $\mathcal{O}^{ss}(\mathbf{H}_\kappa)$ as the full subcategory of $\mathcal{O}(\mathbf{H}_\kappa)$ consisting of $\mathbb{F}[\underline{u}]$ -semisimple modules.

It follows that $\dim M_\zeta < \infty$ for any $\zeta \in \mathbb{F}^n$ and for any $M \in \mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$ or $M \in \mathcal{O}^{ss}(\mathbf{H}_\kappa)$.

Proposition 7.1. *An $\mathbf{H}_\kappa$ -module M is in $\mathcal{O}^{ss}(\mathbf{H}_\kappa)$ if and only if $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is in $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$.*

Proof. Suppose that $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is in $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$. Then $M \cong \mathfrak{N}(\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M) \subseteq \tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$, and it is in $\mathcal{O}^{ss}(\mathbf{H}_\kappa)$.

Suppose that M is in $\mathcal{O}^{ss}(\mathbf{H}_\kappa)$. Since $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M = \mathbb{F}[\pi, \pi^{-1}] \otimes_{\mathbb{F}[\pi]} M$ by Corollary 3.6, any element in $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is of the form $\pi^k v$ for some $k \in \mathbb{Z}$ and $v \in M$. The relation (2.1) implies that if $v \in M$ is a weight vector for $u_1, \dots, u_n$ then $\pi^k v$ is also a weight vector. This implies that $\tilde{\mathbf{H}}_\kappa \otimes_{\mathbf{H}_\kappa} M$ is in $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_\kappa)$. $\square$

Let $m \in \mathbb{Z}_{\geq 1}$. Let $\Lambda^+(m, n)$ denote the set of partitions of n with m nonzero components:

$$\Lambda^+(m, n) = \left\{ \lambda = (\lambda_1, \lambda_2, \dots, \lambda_m) \in \mathbb{Z}^m \mid \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m > 0, \sum_{i \in [1, m]} \lambda_i = n \right\}.$$

For $\lambda \in \Lambda^+(m, n)$, let E_λ denotes the irreducible $\mathfrak{S}_n$ -module corresponding to the partition (or the Young diagram) λ . We write $\mathcal{L}_\kappa(\lambda)$ and $\tilde{\mathcal{L}}_\kappa(\lambda)$ for $\mathcal{L}_\kappa(E_\lambda)$ and $\tilde{\mathcal{L}}_\kappa(E_\lambda)$ respectively.

Let $\text{Irr}(\mathcal{O}^{ss}(\mathbf{H}_\kappa))$ (resp. $\text{Irr}(\mathcal{O}(\tilde{\mathbf{H}}_\kappa))$) denote the set of isomorphism classes of the irreducible modules in $\mathcal{O}^{ss}(\mathbf{H}_\kappa)$ (resp. $\mathcal{O}(\tilde{\mathbf{H}}_\kappa)$). Recall that the assignment $\lambda \mapsto \mathcal{L}_\kappa(\lambda)$ gives a one-to-one correspondence $\bigsqcup_{m \in [1, n]} \Lambda^+(m, n) \xrightarrow{\sim} \text{Irr}(\mathcal{O}(\mathbf{H}_\kappa))$ ([GGOR, Section 2.5.1]).

Now let $\kappa \in \mathbb{Z}$, and set

$$\Lambda_{\kappa}^{+}(m, n) = \{\lambda \in \Lambda^{+}(m, n) \mid \kappa - m - \lambda_1 + \lambda_m \in \mathbb{Z}_{\geq 0}\}.$$

Proposition 7.1 and the classification of the irreducible modules in $\mathcal{O}^{ss}(\tilde{\mathbf{H}}_{\kappa})$ ([Ch3, Theorem 6.5] [SV, Corollary 4.23]) implies the following:

Theorem 7.2. (i) For $\kappa \in \mathbb{Z}_{>0}$, the assignment $\lambda \mapsto \mathcal{L}_{\kappa}(\lambda)$ gives a one-to-one correspondence

$$\bigsqcup_{m \in [1, n]} \Lambda_{\kappa}^{+}(m, n) \xrightarrow{\sim} \text{Irr}(\mathcal{O}^{ss}(\mathbf{H}_{\kappa})).$$

(ii) For $\kappa \in \mathbb{Z}_{<0}$, the assignment $\lambda \mapsto {}^{\sigma}\mathcal{L}_{-\kappa}(\lambda)$ gives a one-to-one correspondence

$$\bigsqcup_{m \in [1, n]} \Lambda_{-\kappa}^{+}(m, n) \xrightarrow{\sim} \text{Irr}(\mathcal{O}^{ss}(\mathbf{H}_{\kappa})),$$

where ${}^{\sigma}\mathcal{L}_{-\kappa}(\lambda)$ denotes the $\mathbf{H}_{\kappa}$ -module given by the composition $\mathbf{H}_{\kappa} \xrightarrow{\sigma} \mathbf{H}_{-\kappa} \rightarrow \text{End}(\mathcal{L}_{-\kappa}(\lambda))$.

Remark 7.3. We stated Theorem 7.2 for $\kappa \in \mathbb{Z}$ since we treated mainly this case in [SV], but most of the results obtained there can be generalized for general $\kappa \in \mathbb{F} \setminus \{0\}$. In particular it follows that Theorem 7.2 can be generalized by setting $\Lambda_{\kappa}^{+}(m, n) = \Lambda^{+}(m, n)$ for $\kappa \in \mathbb{F} \setminus \mathbb{Q}$, and

$$\Lambda_{\kappa}^{+}(m, n) = \{\lambda \in \Lambda^{+}(m, n) \mid r - m - \lambda_1 + \lambda_m \in \mathbb{Z}_{\geq 0}\}$$

for $\kappa = r/p \in \mathbb{Q}_{>0}$ with $r, p \in \mathbb{Z}_{>0}$, $(r, p) = 1$ (cf. [Ch3, Theorem 6.5]).

Remark 7.4. For $\lambda \in \Lambda^{+}(m, n)$, it holds that ${}^{\sigma}\mathcal{L}_{-\kappa}(\lambda) \cong \mathcal{L}_{\kappa}(\lambda')$, where λ' is the conjugate of λ :

$$\lambda'_i = \#\{a \in [1, m] \mid \lambda_a \geq i\} \quad (i \in [1, \lambda_1]).$$

Remark 7.5. It can be seen that the Knizhnik-Zamolodchikov functor investigated in [GGOR] (and in [VV] for $\tilde{\mathbf{H}}_{\kappa}$) transforms the irreducible representations $\mathcal{L}_{\kappa}(\lambda)$ for $\lambda \in \Lambda_{\kappa}^{+}(m, n)$ to Wenzl's representations [We] of the Hecke algebra.

REFERENCES

- [AST] T. Arakawa, T. Suzuki and A. Tsuchiya, *Degenerate double affine Hecke algebras and conformal field theory*, in *Topological Field Theory, Primitive Forms and Related Topics* (Kyoto 1996) (Ed. M Kashiwara, A. Matsuo, K Saito and I. Satake eds.) Progr. Math., vol 160, Birkhäuser, 1998, 1–34.

- [BEG] Y. Berest, P. Etingof and V. Ginzburg, *Finite dimensional representations of rational Cherednik algebras*, Int. Math. Res. Not. **19** (2003), 1053–1088.
- [Ch1] I. V. Cherednik, *A unification of Knizhnik-Zamolodchikov and Dunkl operators via affine Hecke algebras*. Invent. Math. **106** (1991), No. 2, 411–431.
- [Ch2] I. Cherednik, *Double affine Hecke algebras, Knizhnik-Zamolodchikov equations, and Macdonald’s operators*, Int. Math. Res. Not. **9** (1992), 171–179.
- [Ch3] I. V. Cherednik, *Double affine Hecke algebras and differential Fourier transforms*, Invent. Math. **152** (2003), No. 2, 213–303.
- [Ch4] I. V. Cherednik, *Double affine Hecke algebras*, London Mathematical Societies Lecture Note Series, 319, Cambridge University Press (2005).
- [DO] C. Dunkl and E. Opdam, *Dunkl operators for complex reflection groups*, Proc. London Math. Soc. **86** (2003), 70–108.
- [DX] C. Dunkl and Y. Xu, *Orthogonal polynomials of several variables*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2001.
- [EG] P. Etingof and V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism*, Invent. Math. **147** (2002), No. 2, 243–348.
- [Gr] M. G. Green, *The affine q -Schur algebra*, J. Algebra **215** (1999), No. 2, 379–411.
- [GGOR] V. Ginzburg, N. Guay, E. Opdam and R. Rouquier, *On the category $\mathcal{O}$ for rational Cherednik algebras*, Invent. Math. **154** (2003), No. 3, 617–651.
- [Ka] S. Kakei, *Intertwining operators for a degenerate double affine Hecke algebra and multivariable orthogonal polynomials*, Jour. of Math. Phys. **39** (1998), No. 9, 4993–5006.
- [Su] T. Suzuki, *Classification of simple modules over degenerate double affine Hecke algebras of type A* , Int. Math. Res. Not. **43** (2003), 2313–2339.
- [SV] T. Suzuki and M. Vazirani, *Tableaux on periodic skew diagrams and irreducible representations of the double affine Hecke algebra of type A .*, Int. Math. Res. Not. **27** (2005), 1621–1656.
- [Va] E. Vasserot, *Induced and simple modules of double affine Hecke algebras*, Duke Math. J. **126** (2005), no. 2, 251–323.
- [VV] M. Varagnolo and E. Vasserot, *From double affine Hecke algebras to quantized affine Schur algebras*, Int. Math. Res. Not. **26** (2004), 1299–1333.
- [We] H. Wenzl, *Hecke algebras of type A and subfactors* Invent. Math. **92** (1988), No. 2, 349–383.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY,
KYOTO, 606-8502, JAPAN

E-mail address: takeshi@kurims.kyoto-u.ac.jp